

AD-A125 768

MOMENTS OF THE MINIMUM OF A RANDOM WALK AND COMPLETE  
CONVERGENCE(U) STANFORD UNIV CA DEPT OF STATISTICS  
H HOGAN JAN 83 TR-21 N00014-77-C-0306

1/1

UNCLASSIFIED

F/G 12/1


NL

END

FILMED

101

01/8



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

12

MOMENTS OF THE MINIMUM OF A RANDOM WALK  
AND COMPLETE CONVERGENCE

BY

MICHAEL HOGAN

TECHNICAL REPORT NO. 21  
JANUARY 1983

PREPARED UNDER CONTRACT  
N00014-77-C-0306 (NR-042-373)  
FOR THE OFFICE OF NAVAL RESEARCH

Reproduction in Whole or in Part is Permitted  
for any purpose of the United States Government

Approved for public release; distribution unlimited.

ADA 125768

DMC FILE COPY

DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA



DTIC  
ELECTE  
MAR 17 1983  
D

MOMENTS OF THE MINIMUM OF A RANDOM WALK  
AND COMPLETE CONVERGENCE

by

Michael Hogan  
Stanford University

Technical Report No. 21

January 1983

Prepared Under the Auspices  
of  
Office of Naval Research Contract  
N00014-77-C-0306 (NR-042-373)

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

Department of Statistics  
Stanford University  
Stanford, California



MOMENTS OF THE MINIMUM OF A RANDOM WALK  
AND COMPLETE CONVERGENCE

by

Michael Hogan  
Department of Statistics  
Stanford University

Abbreviated Title: Moments of Minimum of a Random Walk

Key Words: Random Walk, Renewal Theorem, Complete Convergence,  
Strong Law of Large Numbers

Summary: Moments of the Minimum of a Random walk and Complete Convergence.

Let  $S_n$  be a random walk with positive drift. Let  $S_{\min} = \inf_{n \geq 0} \{S_n\}$ . New proofs are given of the following: For  $p \geq 1$   $E|S_{\min}|^p < \infty \Leftrightarrow E(|S_1|^{p+1} 1_{(S_1 < 0)}) < \infty$ ;  $\sum P\{S_n < 0\} < \infty \Leftrightarrow E(|S_1|^2 1_{(S_1 < 0)}) < \infty$ , and some related results.

## 1. Introduction

This paper gives new proofs of the equivalences that are stated as Theorem 1 in Section 3. Robbins and Hsu [6] first showed  $c \Rightarrow e$  in 1948 with  $p = 1$ . They considered the problem in the context of a random walk generated by  $X_1$  with  $E X_1 = 0$ , and showed that  $E X_1^2 < \infty \Rightarrow \sum_n P\{\frac{|S_n|}{n} > \varepsilon\} < \infty$ . They called the finiteness of this sum complete convergence. It implies the strong law of large numbers by an application of the Borel-Cantelli lemma. Erdos [3] proved the reverse direction in 1949 and Baum and Katz [1] added the equivalence of (d) in 1965. Kiefer and Wolfowitz [5] established the equivalence of (c) and (f) and the  $(c) \Leftrightarrow (g)$  is in Taylor [7]. Independent discovery of both of the results were credited by the respective authors to unpublished work of Darling, Erdos and Kakutani. These results are partially restated as Theorems 2 and 3 of section 4.

The new proofs provide an  $\varepsilon$ -free approach to these problems. The elementary Renewal Theorem, time reversal, and Wald's identities are the primary tools, and suffice for the case  $p=2$ . For larger  $p$ , the martingale conditional square function has to be used to replace Wald's identities to show the existence of moments in stopped random walks.

## 2. Notation and Conventions.

Fix the following notation and conventions.  $X_1$  is an i.i.d. sequence with  $\mu = EX_1 > 0$ ;  $X^- = -X 1_{\{X < 0\}}$ ;  $S_0 = 0$ ,  $S_i = \sum_{j=1}^i X_j$  for  $i > 0$ ;  $S_{\min} = \inf\{S_i; i > 0\}$ ;  $\tau_+$  is the first strict ascending ladder epoch,  $\tau_+^{(j)}$  is the  $j^{\text{th}}$  strict ascending ladder epoch;  $\tau_-$  is the first weak descending ladder epoch, or  $+\infty$  if none exists,  $\tau_-^{(j)}$  is the  $j^{\text{th}}$  weak descending ladder epoch, or  $+\infty$  if none exists (see Feller [4], Sec. 12.1 for definition);

$$L(0) = \sum_{n=1}^{\infty} 1_{\{\inf_{j \geq n} S_j \leq 0\}}, \quad N(0) = \sum_{n=1}^{\infty} 1_{\{S_n \leq 0\}},$$

i.e.  $L(0)$  is the last time the process is non-positive, and  $N(0)$  is the number of times the process is non-positive;  $\inf\{\} = \infty$ ;  $\tau(a) = \inf\{n > 0: S_n \geq a\}$ ;  $t(a) = \inf\{n > 0: S_n \leq a\}$ ;  $K$  and  $C$  will be positive constants, not necessarily the same from line to line;  $E\{Y; A\} = E(Y1_A)$ ;  $\omega = (X_1, X_2, \dots)$ ,  $\omega_a^+ = (X_{a+1}, X_{a+2}, \dots)$ .  $E^x$  denotes expectation of the random walk started from  $x$ ;  $E = E^0$ .

### 3. Statement and Proof of Theorem:

Theorem 1: For  $p \geq 2$  the following are equivalent:

- a).  $E(\tau_+^p) < \infty$
- b).  $E(\tau_-^{p-1}; \tau_- < \infty) < \infty$
- c).  $E((X^-)^p) < \infty$
- d).  $E(L(0)^{p-1}) < \infty$
- e).  $E(N(0)^{p-1}) < \infty$
- f).  $E(|S_{\min}|^{p-1}) < \infty$
- g).  $E(|S_{\tau_-}|^{p-1}; \tau_- < \infty) < \infty$ .

Four lemmas will be given first, then the proof proceeds as follows:

$a \Leftrightarrow b$ ;  $a \Leftrightarrow c$ ;  $d \Rightarrow b$ ;  $a$  and  $b \Rightarrow d$ ;  $d \Rightarrow e$ ;  $e \Rightarrow g$ ;  $g \Rightarrow f$ ;  $f \Rightarrow e$ ;  
 $f \Rightarrow c$ .

Lemma 1: If  $E(\tau_+^p) < \infty$  then  $E(t(-x)^{p-1}; t(-x) < \infty) < K \quad \forall x > 0$ .

Proof: By time-reversal one has,  $\forall K > 0$ ,

$$\begin{aligned}
 P\{t(-K) = n\} &= P\{S_1 > -K, \dots, S_{n-1} > -K, S_n < -K\} \\
 &= P\{S_n - S_{n-1} > -K, \dots, S_n - S_1 > -K, S_n < -K\} \\
 &\leq P\{S_1 < 0, \dots, S_n < 0\} \\
 &= P\{\tau_+ > n\}.
 \end{aligned}$$

Multiply the first and last lines by  $n^{p-1}$  and summing implies the result, with  $K = C_p E(\tau_+^p)$ .

Lemma 2: If  $E(\tau_+^p) < \infty$  then  $E(\tau(y)^p) \leq K(y+1)^p \quad \forall y > 0$ .

Proof: First notice that  $E(\tau_+^p) < \infty \Rightarrow E(\tau(y)^p) < \infty, \forall y > 0$ . For if  $P\{X < 0\} > 0$  one conditions on the random walk at time 1 to show  $E(\tau(\varepsilon)^p) < \infty, \forall \varepsilon > 0$ , from which  $E(\tau(y)^p) < \infty \quad \forall y$  follows as below, if  $P\{X < 0\} = 0$  the one-sided hitting problem is the same as a two-sided problem, for which Stein's Lemma (cf Feller [5], Sec. 18.2) says  $\tau(y)$  has moments of all orders. To proceed with the proof, observe that for  $K > 0$  an integer

$\tau(Ky) \leq \tau(y) + \tau(y)(\omega_{\tau(y)}^+) + \dots + \tau(y)(\omega_{\tau((K-1)y)}^+)$  and the  $\tau(y)(\omega_{\tau(jy)}^+)$  are i.i.d. Hence by Minkowski's Inequality.

$$E(\tau(Ky)^p) \leq K^p E(\tau(y)^p)$$

and so



$$\begin{aligned}
E(\tau(y)^p) &\leq E \tau([y] + 1)^p \\
&\leq E(\tau(1)^p)([y] + 1)^p \\
&\leq E(\tau(1)^p)(y + 1)^p
\end{aligned}$$

Lemma 3: For  $x > 0$  let

$$\begin{aligned}
R_{-x} &= -S_{t(-x)} - x, \quad t(-x) < \infty \\
&0, \quad t(-x) = \infty.
\end{aligned}$$

Then  $\forall p \geq 1 \ E((X^-)^p) < \infty \Rightarrow E(R_{-x}^{p-1}; t(-x) < \infty) < K$ , where  $K$  is independent of  $x$ .

Proof: This is essentially the same as Theorem 2.4 in Woodroffe [8].

$$\begin{aligned}
P\{R_{-x} > y\} &= \sum_{n=1}^{\infty} P\{t(-x) \geq n, S_n < -x - y\}. \\
&\leq \sum_{n=1}^{\infty} P\{S_{n-1} \geq -x, S_n < -x - y\} \\
&= \sum_{n=1}^{\infty} \int_{-x}^{\infty} F(-x - y - s) F^{*(n-1)}(ds) \\
&= \int_{-x}^{\infty} F(-x - y - s) U(ds) \\
&\leq C \sum_{k > -x} F(-x - y - k) \\
&\leq C \sum_{k > 0} F(-y - k) \\
&\leq C \int_{-\infty}^{-y} F(z) dz.
\end{aligned}$$

where  $F^{*j}$  is the  $j$ -fold convolution of  $F$  with itself, and  $U$  is the renewal measure:  $U(x) - U(y) = \sum_{m=0}^{\infty} [F^{*m}(x) - F^{*m}(y)]$ . Multiplying the first and last statements by  $y^{p-2}$  and integrating gives the stated result.

Lemma 4:  $E(|S_{\tau_-}|^p; \tau_- < \infty) < \infty \Rightarrow E((X^-)^{p+1}) < \infty \forall p > 0$ .

Proof: The statement is invariant under change of scale, so if  $X$  is lattice one may assume that the span of  $X$  is less than 1. It may also be assumed that  $X$  is not bounded below, for otherwise the statement of the lemma is trivial. In this case, with

$$R_x = S_{\tau(x)} - x,$$

since the asymptotic distribution of  $R_x$  has positive mass on  $[0,1)$  (see Woodroffe [8], Sec. 2.2),

$$0 < r = \inf_{x \geq 0} P\{R_x < 1\}.$$

By time-reversal, for  $n \geq 0$

$$\begin{aligned} & P\{S_{\tau_-} \in (-n-1, -n), \tau_- < \infty\} \\ &= \sum_{j=1}^{\infty} P\{S_1 > 0, \dots, S_{j-1} > 0, S_j \in (-n-1, -n)\} \\ &= \sum_{j=1}^{\infty} P\{S_j > S_1, \dots, S_j > S_{j-1}, S_j \in (-n-1, -n)\} \\ &\geq \sum_{j=1}^{\infty} P\{-n-1 > S_1, \dots, -n-1 > S_{j-1}, S_j \in (-n-1, -n)\} \\ &= \sum_{j=1}^{\infty} P\{\tau_{-n-1} = j, R_{-n-1} < 1\} \\ &= P\{R_{-n-1} < 1\}. \end{aligned}$$

And

$$\begin{aligned}
 P\{R_{-n-1} < 1\} &\geq P\{R_{-n-1} < 1, X_1 \leq -n-1\} \\
 &= \int_{n+1}^{\infty} P\{R_{x-n-1} < 1\} P\{X_1^- \in dx\}. \\
 &\geq r P\{X_1^- > n+1\}.
 \end{aligned}$$

Thus

$$P\{S_{\tau_-} \in (-n-1, -n)\} \geq r P\{X_1^- \geq n+1\}.$$

Multiplying by  $n^p$  and summing gives the statement of the lemma.

Proof of the Theorem.

a  $\Rightarrow$  b  $E(\tau_+^p) < \infty \Rightarrow E(\tau_-^{p-1}; \tau_- < \infty) < \infty$ . By a standard time reversal argument (See Feller [4], Sec. 12.2),

$$P\{\tau_+ > n\} = \sum_{j=1}^{\infty} P\{\tau_-^{(j)} = n\}.$$

In particular

$$P\{\tau_+ > n\} \geq P\{\tau_- = n\}.$$

Multiplying by  $n^{p-1}$  and summing gives the result.

b  $\Rightarrow$  a  $E(\tau_-^{p-1}; \tau_- < \infty) < \infty \Rightarrow E(\tau_+^p) < \infty$ . Note that conditioned on  $\tau_-^{(j)} < \infty$   $\tau_-^{(j)} = \sum_{i=1}^j Y_i$ , where the  $Y_i$  are i.i.d. with

$$P\{Y_1 < y\} = P\{\tau_- < y \mid \tau_- < \infty\}. \text{ Thus } \sum_{m=1}^{\infty} n^{p-1} P\{\tau_-^{(j)} = n\} =$$

$$E(|\tau_-^{(j)}|^{p-1}; \tau_-^{(j)} < \infty) \leq j^{p-1} E(\tau_-^{p-1} \mid \tau_- < \infty) P\{\tau_- < \infty\}^j$$

So

$$\sum_{m=1}^{\infty} n^{p-1} P\{\tau_+ > n\} \leq \sum_{j=1}^{\infty} j^{p-1} P\{\tau_- < \infty\}^{j-1} E(\tau_-^{p-1}; \tau_- < \infty)$$

$$\leq K E(\tau_-^{p-1}; \tau_- < \infty).$$

$$\underline{a} \Rightarrow \underline{c} \quad E(\tau_+^p) < \infty \Rightarrow E((X^-)^p) < \infty.$$

By the Elementary Renewal Theorem (Chung [9], Thm. 5.5.2),  $\exists c, K > 0$  such that  $E(\tau(x)) > cx \forall x > K$ . So  $E(\tau(x)^p) \geq (E(\tau(x)))^p \geq c^p x^p \forall x > K$

Conditioning on the first step of the random walk gives

$$\infty > E(\tau_+^p)$$

$$\geq \int_0^{\infty} E(\tau(x))^p P\{X^- \in dx\}.$$

$$\geq \int_K^{\infty} c^p x^p P\{X^- \in dx\}.$$

The last line implies  $E((X^-)^p) < \infty$ .

$$\underline{c} \Rightarrow \underline{a} \quad E((X^-)^p) < \infty \Rightarrow E(\tau_+^p) < \infty. \text{ It suffices to assume } X_i \leq c_x \text{ for some}$$

$c > 0$ ; for,  $X_i$  can be truncated above to give  $\tilde{X}_i$  with  $E \tilde{X}_i > 0$ , and  $\tau_+$  for the random walk generated by the  $\tilde{X}_i$  is larger than that for the  $X_i$  random walk, so if the claim can be proven for the  $\tilde{X}_i$  process it follows for the  $X_i$  process.

In this case it may be assumed that the  $X_i$  have at least 2 moments. Wald's identity for the 2<sup>nd</sup> moment gives

$$E(S_{\tau_+} - \mu \tau_+)^2 = (\text{Var } X_1) E(\tau_+) < \infty.$$

But  $S_{\tau_+} < c$  so  $E S_{\tau_+}^2 < \infty \Rightarrow E \tau_+^2 < \infty$ . Let  $\hat{q} = \sup\{p \geq 2:$

$E((X^-)^q) < \infty \Rightarrow E(\tau_+^q) < \infty \forall 2 \leq q \leq p\}$ . Suppose  $\hat{q} < \infty$ . Let  $\hat{q} \leq q < 2\hat{q}$ .

Then  $E(\tau_+^{q/2}) < \infty$ . Therefore, by Burkholder and Gundy [2], Theorem 5.3.

$$E | S_{\tau_+} - \mu \tau_+ |^q < \infty$$

from which  $E \tau_+^q < \infty$  follows as above. This is a contradiction.

$$\underline{d} \Rightarrow \underline{b} \quad E(L(0)^{p-1}) < \infty \Rightarrow E(\tau_-^{p-1}; \tau_- < \infty) < \infty.$$

Proof:  $L(0) \geq \tau_-^1 \{\tau_- < \infty\}$ .

$$\underline{a} \text{ and } \underline{b} \Rightarrow \underline{d} \quad E(\tau_+^p) < \infty, \text{ and } E(\tau_-^{p-1}; \tau_- < \infty) < \infty \Rightarrow E(L(0)^{p-1}) < \infty.$$

The idea of the proof is to express  $L(0)$  as a sum of successive trips above and below the origin, until the random walk stays permanently above 0. Finding the random walk above 0 one must know the  $p-1^{\text{st}}$  moment of the expected time to get back below 0 must be bounded no matter where the process is, provided it ever does. This is the content of Lemma 1. Having hit below 0 one must know that the  $p-1^{\text{st}}$  moment of the expected time to reach 0 is finite. According to Lemma 2 this quantity is bounded by  $K \int_0^\infty (|y| + 1)^{p-1} F(dy)$ , where  $F$  denotes the hitting place of the non-positive axis.

Lemma 3 provides a uniform bound on the  $p-1^{\text{st}}$  moments of these distributions  $F$ . The proof is then finished by observing that, because of the positive drift, this cycling behavior can only be repeated a few times.

Proof: Set  $p = (1-q) = P\{\tau_- = \infty\}$ . Define

$$T_1 = \inf\{K \geq 1: S_K > 0 \text{ and } \exists m < K \text{ with } S_m \leq 0,$$

and for  $n > 1$

$$T_n = \inf\{K > T_1 + \dots + T_{n-1} = S_K > 0 \text{ and } \exists T_1 + \dots + T_{n-1} < m < K \text{ with } S_m \leq 0\} - (T_1 + \dots + T_{n-1}), T_{n-1} < \infty$$

$$= \infty, T_{n-1} = \infty$$

$$[3.1] \quad |L(0)|^{p-1} \leq \sum_{m=1}^{\infty} |T_1 + \dots + T_n|^{p-1} 1_{\{T_n < \infty, T_{n+1} = \infty\}}.$$

$$E(|T_1 + \dots + T_n|^{p-1}; T_n < \infty, T_{n+1} = \infty)$$

$$\leq E(|T_1 + \dots + T_n|^{p-1}; T_n < \infty)$$

$$\leq n^{p-1} E(T_1^{p-1} + \dots + T_n^{p-1}; T_n < \infty)$$

$E(T_i^{p-1}; T_n < \infty)$  is estimated separately when  $i=n$ , and  $i < n$ . First the case  $i < n$ .

$$E(T_i^{p-1}, T_n < \infty) = E(E(T_i^{p-1}; T_n < \infty | \mathcal{F}_{T_1 + \dots + T_{n-1}}))$$

$$= E(T_i^{p-1}; T_{n-1} < \infty P\{T_n < \infty | \mathcal{F}_{T_1 + \dots + T_{n-1}}\}; T_{n-1} < \infty)$$

$$\leq q E(T_i^{p-1}; T_{n-1} < \infty) \quad (*)$$

and

$$E(T_n^{p-1}; T_n < \infty) = E(E(T_n^{p-1}; T_n < \infty \mid \mathcal{F}_{T_1 + \dots + T_{n-1}}))$$

$$= E(E^{S_{T_1} + \dots + T_{n-1}}(T_1^{p-1}; T_1 < \infty); T_{n-1} < \infty).$$

Consider for  $x > 0$

$$\begin{aligned} E^x\{T_1^{p-1}; T_1 < \infty\} &= E^x\{(\tau_- + \tau(-S_{\tau_-})(\omega_{\tau_-}^+))^{p-1}; \tau_- < \infty\} \\ &\leq 2^{p-1}(E^x(\tau_-^{p-1}; \tau_- < \infty) + E^x(E^{S_{\tau_-}}(\tau(0)^{p-1}; \tau_- < \infty))). \end{aligned}$$

The first term is  $\leq K$  by Lemma 1. For the 2<sup>nd</sup>, using lemmas 2 and 3 it follows that

$$\begin{aligned} &E^x(E^{S_{\tau_-}}(\tau(0)^{p-1}; \tau_- < \infty)) \\ &= \int_{-\infty}^0 E^y(\tau(0)^{p-1}) P^x\{S_{\tau_-} \in dy, \tau_- < \infty\} \\ &\leq K' \int_{-\infty}^0 |y + 1|^{p-1} P^x\{S_{\tau_-} \in dy, \tau_- < \infty\}. \\ &\leq K''. \end{aligned}$$

Thus

$$\begin{aligned} E(T_n^{p-1}; T_n < \infty) &\leq K P\{T_{n-1} < \infty\} \\ &\leq K q^{n-1}. (**). \end{aligned}$$

$$\text{Set } a_n = E(T_1^{p-1} + \dots + T_n^{p-1}; T_n < \infty).$$

Summing (\*) from 1 to  $n-1$  and adding (\*\*) gives

$$a_n \leq q a_{n-1} + K q^{n-1},$$

Therefore,  $a_n$  is geometrically decreasing, and  $\sum a_n < \infty$ . A look at 3.1 shows that  $\sum a_n < \infty \Rightarrow E L(0)^{p-1} < \infty$ .

$$\underline{d} \Rightarrow \underline{e} \quad E(L(0)^{p-1}) < \infty \Rightarrow E(N(0)^{p-1}) < \infty$$

Proof:  $L(0) \geq N(0)$ .

$$\underline{e} \Rightarrow \underline{g} \quad E(N(0)^{p-1}) < \infty \Rightarrow E(|S_{\tau_-}|^{p-1}; \tau_- < \infty) < \infty$$

Proof: The amount of time spent getting back above  $\phi$  after having hit below  $\phi$  for the first time is  $\tau(0)(\omega_{\tau_-}^+) 1_{\{\tau_- < \infty\}}$ . So

$$(1 + N(0))^{p-1} \geq \tau(0)(\omega_{\tau_-}^+) 1_{\{\tau_- < \infty\}}.$$

$$\text{and } \infty > E((1 + N(0))^{p-1}) \geq E(\tau^{p-1}(0)(\omega_{\tau_-}^+); \tau_- < \infty)$$

$$= E(E(\tau^{p-1}(0)(\omega_{\tau_-}^+); \tau_- < \infty | F_{\tau_-}))$$

$$= E(E^{S_{\tau_-}}(\tau(0)^{p-1}; \tau_- < \infty) \Rightarrow E(|S_{\tau_-}|^{p-1}; \tau_- < \infty) < \infty$$

as in the last part of a and  $b \Rightarrow d$ .

$$\underline{g} \Rightarrow \underline{f} \quad E(|S_{\tau_-}|^{p-1}; \tau_- < \infty) < \infty \Rightarrow E(|S_{\min}|^{p-1} < \infty)$$

Proof:  $S_{\min}$  can be written as  $Z_n$ , where  $Z_1$  is a random walk with  $P\{Z_1 < y\} = P\{S_{\tau_-} < y\} | \tau_- < \infty$ ,  $P\{M = n\} = P\{\tau_- < \infty\}^n P\{\tau_- = \infty\}$ ,  $n=0, 1, \dots$ , and  $M$  is independent of the  $Z_1$ . This can be seen intuitively by considering the decreasing ladder process, or a quick proof can be based on a comparison of the characteristic functions given in Feller [4], Chapt. 18.  $E(|S_{\tau_-}|^{p-1}; \tau_- < \infty) < \infty \Rightarrow E(|Z_1|^{p-1}) < \infty$ , so

$$\begin{aligned} E(|S_{\min}|^{p-1}) &= \sum_n E(|Z_n|^{p-1}) P\{M = n\} \\ &\leq E(|Z_1|^{p-1}) \sum_n n^{p-1} P\{M = n\} \\ &< \infty. \end{aligned}$$



$$\underline{f \Rightarrow g} \quad E(|S_{\min}|^{p-1}) < \infty \Rightarrow E(|S_{\tau_-}|^{p-1}; \tau_- < \infty) < \infty$$

Proof:  $|S_{\tau_-}| 1_{\{\tau_- < \infty\}} \leq |S_{\min}|$

$$\underline{f \Rightarrow e} \quad E(|S_{\min}|^{p-1}) < \infty \Rightarrow E(N(0)^{p-1}) < \infty.$$

Proof: Since  $E(|S_{\min}|^{p-1}) < \infty$ , then  $E(|S_{\tau_-}|^{p-1}; \tau_- < \infty) < \infty$ . From lemma 4  $E((X^-)^{p+1}) < \infty$  so the result follows from  $c \Rightarrow d \Rightarrow e$ .

$$\underline{f \Rightarrow c} \quad E(|S_{\min}|^{p-1}) < \infty \Rightarrow E((X^-)^p) < \infty$$

Proof: Follows from  $f \Rightarrow g$  and Lemma 4.

#### 4. Remarks and Applications.

Let  $Y_i$  be a i.i.d. sequence of random variables with  $EY_i = 0$ .

Let  $S_n = Y_1 + \dots + Y_n$

$$L(\varepsilon) = \sum_{n=1}^{\infty} 1_{\{\sup_{j \geq n} \frac{|S_j|}{j} > \varepsilon\}},$$

$$N(\varepsilon) = \sum_{n=1}^{\infty} 1_{\{|\frac{S_n}{n}| > \varepsilon\}}.$$

Theorem 2: For  $p \geq 2$

$$(1) \quad E((N(\varepsilon))^{p-1}) < \infty \Leftrightarrow E(|Y|^p) < \infty$$

$$(2) \quad E(L(\varepsilon)^{p-1}) < \infty \Leftrightarrow E(|Y|^p) < \infty$$

$$(3) \quad \sum_{n=1}^{\infty} P\{\sup_{j \geq n} \frac{|S_j|}{j} > \varepsilon\} \cdot n^{p-2} < \infty \Leftrightarrow E|Y|^p < \infty$$

$$(4) \quad \sum_{n=1}^{\infty} P\{|\frac{S_n}{n}| > \varepsilon\} \Leftrightarrow E|Y|^2 < \infty.$$

Remarks: The "only if" part of (4), for  $p=1$  is due to Robbins and Hsu [6], (4) with  $p=1$  is due to Erdos [3], (3) was first proved by Baum and Katz and can be found in [1].

Proof: (1) and (2) follow from the equivalence of c,d, and e by considering the random walks  $S_n \pm n\epsilon$ . (3) is the same as (2) plus the observation that  $P\{L(\epsilon) > n\} = P\{\sup_{j \geq n} \frac{|S_j|}{j} > \epsilon\}$ , and (4) follows similarly from (1).

Let  $X_i$  be i.i.d. random variables with  $E X_i = \mu \in (0, \infty)$ ,  
 $S_n = X_1 + \dots + X_n$ .

Theorem 3: For  $p \geq 1$  the following are equivalent:

- (1)  $E((X^-)^{p+1}) < \infty$
- (2)  $E(|S_{\tau_-}|^p; \tau_- < \infty) < \infty$
- (3)  $E(|S_{\min}|^p) < \infty$ .

Remark: The equivalence of (2) and (3) for  $p=1$  is credited by Taylor [7], to unpublished work of Darling, Erdos and Kakutani, and Taylor adds a proof of the equivalence of (1). Kiefer and Wolfowitz [5] also credit the equivalence of (1) and (3) to unpublished results of Darling, Erdos and Kalutani; and they give their own proof. The moments of the minimum are of interest because the minimum has the distribution of the stationary distribution of a type of queueing process. See [4] p. 198.

Proof: This is the equivalence above, however, the tortuous path via the implications of Theorem 1 can be replaced by lemma 4.

I would like to thank Professor Siegmund for help received on this problem. In particular he showed me the time-reversal proof of  $b \Rightarrow a$ .

## Reference

1. Baum L.E. and Katz M. (1965) Convergence Rates in the Law of Large Numbers. Transactions of Ann. Math. Soc. 120, pp. 108-123.
2. Burkholder, D.L. and Gundy, R.F. (1970) Extrapolation and Interpolation at Qausi-Linear Operators on Martingales Acta Math 124, pp. 249-304.
3. Erdos, P. (1949), On a Theorem of Hsu and Robbins, Ann. Math. Stat., 20, pp. 286-291.
4. Feller, W. (1966), An Introduction to Probability Theory and Its Applications II, 2nd Ed., John Wiley, New York.
5. Kiefer, J. and Wolfowitz, J. (1956) The General Queueing Process, Ann. Math. Stat., 27, pp. 147-161.
6. Robbins H. and Hsu., (1947), Complete Convergence and the Law of Large Numbers, Proc. Nat. Acad. Sciences (33) pp. 25-31.
7. Taylor, H.M. (1972), Bounds for Stopped Partial Sums, Ann. Math. Stat. 43, pp. 733-747.
8. Woodroffe, M. (1981), Non Linear Renewal Theory in Sequential Analysis, University of Michigan, Technical Report No.106.
9. Chung, K. L. (1974) A Course in Probability Theory, 2nd ed., Academic Press, New York.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 21	2. GOVT ACCESSION NO. A125 768	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Moments of the Minimum of a Random Walk and Complete Convergence		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL REPORT
7. AUTHOR(s) Michael Hogan		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Stanford University Stanford, California		8. CONTRACT OR GRANT NUMBER(s) N00014-77-C-0306
11. CONTROLLING OFFICE NAME AND ADDRESS Statistics & Probability Program Code (411 (SP)) Office of Naval Research Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-042-373
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE January 1983
		13. NUMBER OF PAGES 14
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release: Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Random Walk, Renewal Theorem, Complete Convergence, Strong Law of Large Numbers		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let $S_n$ be a random walk with positive drift. Let $S_{\min} = \inf_{n>0} \{S_n\}$ . New proofs are given of the following: For $p \geq 1$ $E S_{\min} ^p < \infty \iff$ $E( S_1 ^{p+1} 1_{(S_1 < 0)}) < \infty$ ; $\sum P\{S_n < 0\} < \infty \iff E( S_1 ^2 1_{(S_1 < 0)}) < \infty$ , and some related results.		

DD FORM 1473 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

3-8

DT